



(ICRP No. 103). By the way, the most accurate volume and dose data is for mammography because of the 100% data collection mandated by MQSA. The volume of mammography was reported in 2006 at 34.4 million mammograms with a population of women (45 yr and older) of 58.2 million. Also included here are data regarding the use of digital breast tomosynthesis.

For Computed Tomography, the American College of Radiology Dose Index Registry (DIR) has been of value to the authors of this report. The compilation of computed tomography dose indices and dose-length product values along with the anatomical sites provides a rich set of data with narrowing error bars. This is important because CT is still the highest medical source of radiation and was essentially unchanged from 2006 to 2016.

At this point, the patient reader will know that we are just getting to the good part – the rest of you are asking “Why did the numbers decrease? Please tell!” The answer is a combination of technology and practice patterns, with the latter being the larger part of the decrease. While the number of CT studies increased by 25% (and population only increased by about 8%), the use of dose reduction methods kept the annual individual effective dose at 1.5 mSv (1.4 mSv in 2006). For other modalities, there were substantial decreases in the number of procedures – this included general and cardiovascular nuclear medicine – which resulted in a 68% decrease in the per individual dose. As the second largest contributor to collective effective dose, this was a significant effect. However, decreases in the number of chest, abdomen, pelvis and urologic exams resulted in a decrease of about 25%.

Lastly, it is important to recognize that the principle value identified in this report (Effective Dose per individual in the U.S. Population) is obtained by dividing the collective effective dose by the U.S. population, whether exposed or not. Since many people do not get any x-ray, CT or nuclear studies in the course of a year and many get multiple studies. Therefore, this number should not be used in a general epidemiological application, for example, how many cancers will result from medical imaging this year? The value of this data point is the historical comparison and trends. As such, it helps us to understand the effect of an aging population, Medicare reimbursement, technology and medical practice patterns have on the use of imaging. It is also still useful as a way to review overall population health. As health care professionals, we can say (informed as we are) that the use of medical imaging contributes roughly the same radiation exposure to the U.S. population as radon, cosmic radiation, consumer products and nuclear power combined.

By the way, the National Council on Radiation Protection and Measurements was chartered by the U.S. Congress and seeks to formulate, disseminate guidance and recommendations on radiation protection and measurements. Your friendly neighborhood medical physicist relies on the activities and publications of this organization for a variety of matters – our peers are often quite involved in these.

However, they also offer guidance to government entities like NASA (spaceflight dosimetry and radiobiology) and federal and state regulatory agencies. The reports are collaborative engagements, often involving twenty or more experts in the field.

